



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 211/18, 409/06, 211/70, 295/135, 317/58, 211/52, 211/10, 401/12, 409/14, 401/04, 401/10, A61K 31/445, 31/50</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/33721 (43) International Publication Date: 14 December 1995 (14.12.95)</p>
<p>(21) International Application Number: PCT/DK95/00230 (22) International Filing Date: 8 June 1995 (08.06.95) (30) Priority Data: 0649/94 8 June 1994 (08.06.94) DK (71) Applicant (for all designated States except US): H. LUND- BECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Copenhagen- Valby (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): PERREGAARD, Jens, Kristian [DK/DK]; Thyrasvej 22, DK-3630 Jægerspris (DK). STENBERG, John, Willie [DK/DK]; Vesterbrogade 55A, DK-1620 Copenhagen (DK). HANSEN, Bitten [DK/DK]; Tørnergården 68, DK-4600 Køge (DK). (74) Agent: MEIDAHN PETERSEN, John; H. Lundbeck a/s, Patent Dept., Ottiliavej 9, DK-2500 Copenhagen-Valby (DK).</p>		<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i></p>
<p>(54) Title: 4-ARYL-1-(INDANMETHYL, DIHYDROBENZOFURANMETHYL OR DIHYDROBENZOTHIOPHEN- EMETHYL)PIPERIDINES, TETRAHYDROPYRIDINES OR PIPERAZINES</p> <div style="text-align: center;"> <p>(I)</p> </div> <p>(57) Abstract</p> <p>4-Aryl-1-(indanmethyl, dihydrobenzofuranmethyl or dihydrobenzothiophenemethyl)piperidine, -tetrahydropyridine or -piperazine compounds of general formula (I), wherein one of X and Y is CH₂ and the other one is CH₂, O or S; Z is N, C, CH or COH; Ar is an optionally substituted aryl group; R¹ is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, acyl, thioacyl, alkylsulfonyl, trifluoromethylsulfonyl, arylsulfonyl, a group R⁹VCO- where V is O or S and R⁹ is alkyl or aryl, or a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- wherein R¹⁰ and R¹¹ are hydrogen, alkyl or aryl, or R¹⁰ and R¹¹ are linked to form a ring; R² is hydrogen, alkyl, cycloalkyl or cycloalkylalkyl; or R¹ and R² are linked to form a ring; R³ - R⁵ are hydrogen, halogen, alkyl, alkylcarbonyl, phenylcarbonyl, alkoxy, alkylthio, hydroxy, alkylsulfonyl, cyano, trifluoromethyl, cycloalkyl, cycloalkylalkyl or nitro; R⁶ and R⁷ are hydrogen or alkyl or they are linked to constitute 3-7-membered ring; R⁸ is hydrogen or alkyl; have effects at central serotonergic receptors and are therefore useful in the treatment of certain psychic and neurologic disorders.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

4-Aryl-1-(indanmethyl, dihydrobenzofuranmethyl or dihydrobenzothiophenemethyl)piperidines, tetrahydropyridines or piperazines.

Field of the Invention

5 The present invention relates to a novel class of 4-aryl-1-(indanmethyl, dihydrobenzofuranmethyl or dihydrobenzothiophenemethyl)piperidine, -tetrahydropyridine or -piperazine compounds having effects at central serotonergic receptors. These methylamine compounds are therefore useful in the treatment of certain psychic
10 and neurologic disorders.

Background of the Invention.

A few aminomethylindan, -dihydrobenzofurane and -dihydrobenzothiophene com-
15 pounds are known from the prior art.

So, EP patent 0 281 261 discloses 1-aminomethylindan, 3-aminomethylbenzofurane and 3-aminomethylbenzothiophene derivatives with a hydroxy group or a substituted hydroxy group in the 6-position (indan) or 5-position (benzofurane,
20 benzothiophene). These compounds were found to show central dopamine agonist activity, in particular to show effect at presynaptic dopamine receptors.

In U.S. Patent No. 4,500,543 certain 1-aminomethylphtalane compounds are said to show adrenergic effects and, accordingly, antihypertensive and heart rate decreasing properties. Said patent generically covers compounds having substituents in
25 the 5-, 6- and/or 7-position.

EP 0325963 A1 discloses among other compounds a class of 1-aminomethyl indan compounds in which the aminomethyl group may constitute a 1-pyrrolylmethyl
30 group which is substituted with thienyl or phenyl. The compounds are claimed to be α_2 antagonists useful in the treatment of depression, metabolic disorders, glaucoma, migraine and hypertension.

Furthermore, EP 0490772 A1 describes i.a. a class of 4-benzofuranyl- or 4-benzodioxanyl-1-indanylmethyl piperazine compounds being 5-HT_{1A} ligands.

EP 0428437 generically covers a very broad class of 1,2-benzisoxazole compounds including certain 3-[1-[(1-indanyl)methyl]-1,2-benzisoxazoles. However, only one such compound is exemplified and in that case without giving any data. The compounds are said to show dopamine and serotonin antagonistic activities.

US Patent No 3,886,168 relates to 1-[(indan-1-yl)methyl]piperidine compounds having antihypertensive activity.

Various effects are known with respect to compounds which are ligands at the different serotonin receptor subtypes. As regards the 5-HT_{2A} receptor, which was previously referred to as the 5-HT₂ receptor, the following effects have e.g. been reported:

The 5-HT_{2A} antagonist ritanserin (Meert, T. F.; Janssen, P. A. J. *Drug. Dev. Res.* **1989**, *18*, 119.) has been shown to be effective in the treatment of anxiety and depression presumably through improvement of the sleep quality. Furthermore, selective, centrally acting 5-HT_{2A} antagonists have been shown to have an effect towards the negative symptoms of schizophrenia and to reduce extrapyramidal side-effects caused by treatment with classical neuroleptics in schizophrenic patients (Gelders, Y.G., *British J. Psychiatry*, **1989**, *155* (suppl.5), 33). Finally, selective 5-HT_{2A} antagonists could be effective in the prophylaxis and treatment of migraine since it is known that 5-HT is involved in migraine attacks. The links between 5-HT and migraine attacks are several and they suggest a number of mechanisms whereby 5-HT may be involved (Scrip Report; "Migraine – Current trends in research and treatment"; PJB Publications Ltd.; May 1991).

The serotonin 5-HT_{2A} antagonist, MDL 100,907 (Sorensen, S.M. et al., *J. Pharmacol. Exp. Ther.* **1993**, *266*, 684-691), and certain compounds within series of 1-phenylindoles (WO 93/12790) and 3-phenylindole derivatives (WO 93/14758) have shown anti-psychotic activity in animal models with indication of no liability to caus

extrapyramidal side effects (EPS).

Clinical studies of known 5-HT_{1A} partial agonists such as e.g. buspirone, 8-[4-[4-(2-pyrimidyl)-1-piperazinyl]butyl]-8-azaspiro[4,5]decane-7,9-dione, gepirone, 4,4-dimethyl-1-[4-[4-(2-pyrimidyl)-1-piperazinyl]butyl]-2,6-piperidinedione, and ipsapirone, 2-[4-[4-(2-pyrimidyl)-1-piperazinyl]butyl]-1,2-benzothiazol-3(2H)-one-1,1-dioxide, have shown that 5-HT_{1A} partial agonists are useful in the treatment of anxiety disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder (Glitz, D. A., Pohl, R., *Drugs* **1991**, *41*, 11). Preclinical studies indicate that also full agonists are useful in the treatment of the above mentioned anxiety related disorders (Schipper, *Human Psychopharmacol.*, **1991**, *6*, S53).

There is also evidence, both clinical and preclinical, in support of the beneficial effect of 5-HT_{1A} partial agonists in the treatment of depression, impulse control disorders and alcohol abuse (van Hest, *Psychopharmacol.*, **1992**, *107*, 474; Schipper et al, *Human Psychopharmacol.*, **1991**, *6*, S53; Cervo et al, *Eur. J. Pharm.*, **1988**, *158*, 53; Glitz and Poh, *Drugs* **1991**, *41*, 11; Grof et al., *Int. Clin. Psychopharmacol.* **1993**, *8*, 167-172; Ansseau et al., *Human Psychopharmacol.* **1993**, *8*, 279-283).

5-HT_{1A} agonists and partial agonists inhibit isolation-induced aggression in male mice indicating that these compounds are useful in the treatment of aggression (Sánchez et al., *Psychopharmacology*, **1993**, *110*, 53-59).

Furthermore, 5-HT_{1A} ligands have been reported to show antipsychotic effect in animal models (Wadenberg and Ahlenius, *J. Neural. Transm.*, **1991**, *83*, 43; Ahlenius, *Pharmacol.&Toxicol.*, **1989**, *64*, 3; Lowe et al., *J. Med. Chem.*, **1991**, *34*, 1860; New et al., *J. Med. Chem.*, **1989**, *32*, 1147; and Martin et al., *J. Med. Chem.*, **1989**, *32*, 1052).

Recent studies also indicate that 5-HT_{1A} receptors are important in the serotonergic modulation of haloperidol-induced catalepsy (Hicks, *Life Science* **1990**, *47*, 1609, Wadenberg et al. *Pharmacol.Biochem. & Behav.* **1994**, *47*, 509-513) suggesting that 5-HT_{1A} agonists are useful in the treatment of EPS induced by conventional

antipsychotic agents such as haloperidol.

5-HT_{1A} agonists have shown neuroprotective properties in rodent models of focal and global cerebral ischaemia and may, therefore, be useful in the treatment of
5 ischaemic disease states (Prehn, *Eur. J. Pharm.* **1991**, *203*, 213).

Pharmacological studies have been presented which indicates that 5-HT_{1A} antagonists are useful in the treatment of senile dementia (Bowen et al, *Trends Neur. Sci.* **1992**, *15*, 84).

10

Both in animal models and in clinical trials it has been shown that 5-HT_{1A} agonists exert antihypertensive effects via a central mechanism (Saxena and Villalón, *Trends Pharm. Sci.* **1990**, *11*, 95; Gillis et al, *J. Pharm. Exp. Ther.* **1989**, *248*, 851). 5-HT_{1A} ligands may, therefore, be beneficial in the treatment of cardiovascular
15 disorders.


5-HT reuptake inhibitors are well known antidepressant drugs.

As 5-HT_{1A} and 5-HT_{2A} receptor ligand classes of compounds and 5-HT reuptake
20 inhibitors have different activities in different animal models predictive of anxiolytic and antiaggressive effects (Perregaard et al., Recent Developments in Anxiolytics. *Current Opinion in Therapeutic Patents* **1993**, *1*, 101-128) and/or in models predictive of effects in other psychic disorders it might also be highly beneficial to treat complex states of anxiety, depression, or other psychic disorders with a drug
25 which have combined serotonergic effects.

Summary of the Invention

It has now been found that certain novel 4-aryl-1-(indanmethyl, dihydrobenzofuran-
30 methyl or dihydrobenzothiophenemethyl)piperidines, -tetrahydropyridines or -piperazines interact potently with central serotonergic receptors, in particular with the 5-HT_{1A} and/or the 5-HT_{2A} receptors.

Accordingly, the present invention relates to novel compounds of the formula I.



- 5 wherein one of X and Y is CH₂ and the other one is selected from the group consisting of CH₂, O, and S;

the dotted line, emanating from Z, indicates an optional bond; when it does not indicate a bond Z is N, CH or COH; and when it indicates a bond Z is C;

10

Ar is phenyl, 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-pyrimidyl, 1-indolyl, 2-indolyl, 3-indolyl, 1-indol-2-onyl, 3-indol-2-onyl, 2- or 3-benzofuranyl, 2- or 3-benzothiophenyl, 1-naphthyl or 2-naphthyl, each optionally substituted with halogen, lower alkyl, lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl, 15 trifluoromethylsulfonyloxy, cycloalkyl, cycloalkyl-lower-alkyl, nitro, amino, lower alkylamino, di-lower alkylamino, acylamino or C₁₋₂ alkylenedioxy;


R₁ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl, cycloalk(en)yl, cycloalk(en)-yl-lower alk(en/yn)yl, aryl, aryl-lower alkyl, acyl, thioacyl, lower alkylsulfonyl, trifluoro- 20 methylsulfonyl, arylsulfonyl,

R₁ is a group R⁹VCO- where V is O or S and R⁹ is lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl, or

R₁ is a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- wherein R¹⁰ and R¹¹ are independently hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl, or R¹⁰ and R¹¹ 25 together with the N-atom to which they are linked, form a pyrrolidinyl, piperidinyl or perhydroazepin group;

R₂ is hydrogen, lower alkyl, cycloalkyl or cycloalkyl-lower-alkyl;

or R¹ and R² together with the N-atom to which they are linked form a group ,



wherein Q is C=O, C=S or CH₂; T is NH, S, O or CH₂; and m is 1-4, inclusive;

- 5 R³ - R⁵ are independently hydrogen, halogen, lower alkyl, lower alkylcarbonyl, phenylcarbonyl, halogen substituted phenylcarbonyl, lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl, cycloalkyl, cycloalkyl-lower-alkyl or nitro;
- 10 R⁶ and R⁷ are each hydrogen or lower alkyl or they are linked together to constitute a 3 - 7-membered carbocyclic ring;

R⁸ is hydrogen or lower alkyl;

- 15 any alkyl, cycloalkyl or cycloalkylalkyl group present being optionally substituted with one or two hydroxy groups, which again are optionally esterified with an aliphatic or aromatic carboxylic acid; and any aryl substituent present being optionally substituted with halogen, lower alkyl, lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl, trifluoromethylsulfonyloxy,
- 20 cycloalkyl, cycloalkyl-lower-alkyl or nitro;

and pharmaceutically acceptable acid addition salts thereof.

The compounds of the invention have been found to show potent affinity to 5-HT_{1A} receptors and/or to 5-HT_{2A} receptors. In addition to the effect at these receptor subtypes, certain of the present compounds also show 5-HT reuptake inhibiting effect.

Accordingly, the compounds of the invention are considered useful in the treatment of positive and negative symptoms of schizophrenia, other psychoses, anxiety disorders, such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, alcohol abuse, impuls control disorders, aggres-

sion, side effects induced by conventional antipsychotic agents, ischaemic disease states, migraine, senile dementia and cardiovascular disorders and in the improvement of sleep.

- 5 In another aspect the invention provides a pharmaceutical composition comprising at least one compound of Formula I as defined above or a pharmaceutically acceptable acid addition salt thereof or prodrug thereof in a therapeutically effective amount and in combination with one or more pharmaceutically acceptable carriers or diluents.

10

In a further aspect the present invention provides the use of a compound of Formula I as defined above or an acid addition salt or prodrug thereof for the manufacture of a pharmaceutical preparation for the treatment of the above mentioned disorders.

15

Detailed Description of the Invention

Compounds of general Formula I exist as optical isomers thereof and such optical isomers are also embraced by the invention.

20

Prodrugs of the compounds of general Formula I are also embraced by the invention.

The term cycloalkyl designates a carbocyclic ring having 3-8 carbon atoms, inclusive, or a bicyclic or tricyclic carbocycle, such as adamantyl.

25

The term lower alkyl refers to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl and 2-methyl-1-propyl. The terms lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkylamino, lower alkylcarbonyl, etc. designate such groups in which the alkyl group is lower alkyl as defined above. Similarly, lower alkenyl and alkynyl, respectively, designate such groups having from two to six carbon atoms, inclusive. Preferred groups are those having up to four carbon

30

atoms.

The term aryl refers to a mono- or bicyclic carbocyclic or heterocyclic aromatic group, such as phenyl, indolyl, thienyl, pyrimidyl, oxazolyl, isoxazolyl, thiazolyl, 5 isothiazolyl, imidazolyl, benzofuranyl, benzothienyl, pyridyl, naphthyl and furanyl, in particular phenyl, pyrimidyl, indolyl, and thienyl.

Halogen means fluoro, chloro, bromo or iodo.

10 As used herein the term acyl refers to a formyl, lower alk(en/yn)ylcarbonyl, aryl-carbonyl, aryl-lower alk(en/yn)ylcarbonyl, cycloalkylcarbonyl, or cycloalkyl-lower-alk(en/yn)ylcarbonyl group.

The term thioacyl is the corresponding acyl group in which the carbonyl group is 15 replaced with a thiocarbonyl group.

The expression alk(en/yn)yl means that the group may be an alkyl, alkenyl, or alkynyl group.

20 In Formula I, X is preferably CH₂ or S and Y is preferably CH₂ and most preferably they are both CH₂.

R¹ is preferably acyl, lower alkyl, lower alkoxy, a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- wherein R¹⁰ is hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl and 25 R¹¹ is hydrogen or lower alkyl or R¹⁰ and R¹¹ together with the N-atom to which they are linked, form a pyrrolidiny, piperidiny or perhydroazepin group. Most preferably, R¹ is formyl, acetyl, methylaminocarbonyl, methylaminothiocarbonyl, dimethylaminocarbonyl, dimethylaminothiocarbonyl, methylsulfonyl, aminocarbonyl, cyclopropylcarbonyl, methyl, pyrrolidinylcarbonyl or 4-fluorophenylaminocarbonyl. 30 R² is preferably hydrogen or lower alkyl, most preferably hydrogen or methyl, or R¹ and R² are linked together to form a 5-7 membered unsubstituted lactam ring or a pyrrolidiny, piperidiny or perhydroazepin.

R³-R⁵ are preferably hydrogen, fluoro, chloro, bromo, methyl, trifluoromethyl or acetyl and R⁶-R⁸ are preferably all hydrogen.

Finally, Ar is preferably phenyl, 3-indolyl, 1-indolyl, or pyrimidyl or phenyl, 3-indolyl,
5 1-indolyl or pyrimidyl substituted with halogen.

A preferred subclass of compounds are those wherein R¹ is acetyl and R² is H and in particular such compounds wherein Ar is indolyl or phenyl substituted with halogen, especially chloro. If Ar is 3-indolyl it is preferably substituted in the 6-
10 position and if it is phenyl, it is preferably substituted in the 4-position.

Another preferred subclass of compounds of the invention are those wherein R¹ is a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- wherein R¹⁰ is hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl and R¹¹ is hydrogen or lower alkyl and R² is hydrogen.

15

In a further preferred subclass of compounds R¹ is hydrogen, lower alkyl or lower alkylsulfonyl in particular methyl or methylsulfonyl and R² is hydrogen or lower alkyl, in particular methyl, or R¹ and R² are linked together to form a pyrrolidinon ring or a pyrrolidinyl ring.

20

Preferred compounds are :

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

(+)-1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

(-)-1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine

25 1-(6-Acetylamino-5-fluoroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-4-fluoroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-4-bromoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-4-nitroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-4-cyanoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

30 1-(6-Acetylamino-5-chloroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-5-bromoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-5-cyanoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

1-(6-Acetylamino-7-chloroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

- 1-(6-Acetylamino-7-fluoroindan-1-ylmethyl)-4-(4-fluorophenyl)piperidin .
1-(5-Acetyl-6-acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
1-(6-Acetylamino-1-methylindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-fluorophenyl)piperidine.
5 1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-fluorophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-methylphenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-dimethylaminophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-aminophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-trifluoromethylphenyl)piperidine.
10 1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperazine.
1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-4-(4-fluorophenyl)piperidine.
1-(6-Acetylamino-1,3-dihydroisobenzofuran-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
15 1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-chlorophenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-chlorophenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-chlorophenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-trifluoromethylsulfonyloxyphenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-dichlorophenyl)piperazine.
20 1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-dichlorophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-methoxyphenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-pyrimidyl)piperidine.
25 1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-pyrimidyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-pyridinyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-thienyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-thienyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-thienyl)piperazine.
30 1-(6-Acetylaminoindan-1-ylmethyl)-4-(1-naphthyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-naphthyl)piperidine.
1-(6-butanoylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine
1-(6-Formylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.

- 1-(6-Formylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperazine.
4-(4-Fluorophenyl)-1-(6-methansulfonylaminoindan-1-ylmethyl)piperidine.
1-(6-Cyclopropylcarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
1-(6-Cyclopentylcarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
5 4-(4-Fluorophenyl)-1-(6-methylaminocarbonylaminoindan-1-ylmethyl)piperidine.
1-[6-(4-Fluorophenyl)aminocarbonylaminoindan-1-ylmethyl]-4-(4-fluorophenyl)piperidine.
4-(4-Fluorophenyl)-1-(6-methylaminothiocarbonylaminoindan-1-ylmethyl)piperidine.
1-(6-Dimethylaminocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
10 1-(6-Dimethylaminothiocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
4-(4-Fluorophenyl)-1-[6-(1-pyrrolidinyl)carbonylaminoindan-1-ylmethyl]piperidine.
1-(6-Aminocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
1-(6-Ethoxycarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine.
15 1-[6-(N,N-dimethylamino)indan-1-ylmethyl]-4-(4-fluorophenyl)piperidine.
3-[1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)piperidin-4-yl]-5-chloro-1 *H*-indole.
3-[1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1 *H*-indole.
20 3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-5-fluoro-1 *H*-indole.
3-[1-(6-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-fluoro-1 *H*-indole.
3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1 *H*-indole.
25 3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1 *H*-indole.
3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1-methyl-1 *H*-indole.
1-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1 *H*-indole.
1-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1 *H*-indole.
3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1 *H*-
30 indole.
4-(4-Fluorophenyl)-1-[6-(1-pyrrolidin-2-onyl)indan-1-ylmethyl]piperidine.
4-(4-Fluorophenyl)-1-[6-(1-piperidin-2-onyl)indan-1-ylmethyl]piperidine.
1-[6-(4-Fluorophenylamino)indan-1-ylmethyl]-4-(4-fluorophenyl)piperidine

- 3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chlorobenzothiophene.
3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chlorobenzothiophene.
3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-5-chlorobenzothiophene.
- 5 2-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chlorobenzothiophene.
3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chlorobenzofurane.
3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1*H*-indol-2-one
3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1-methyl-1*H*-indol-2-one.
- 10 3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1-methyl-1*H*-indol-2-one.
2-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1*H*-indole.
1-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indol-2-one.
3-[1-(6-Methylaminokarbonylaminoindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-
- 15 indole.
3-[1-(6-Methylaminokarbonylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1*H*-indole.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-bromophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-hydroxy-4-(4-chlorophenyl)piperidine.
- 20 1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-trifluoromethyl-4-chlorophenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-chloro-3-thienyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-chloro-2-thienyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperazine.
- 25 1-(6-Methylaminocarbonylaminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperazine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-hydroxy-4-(3-trifluoromethyl-4-chlorophenyl)piperidine.
1-(6-Acetylaminoindan-1-ylmethyl)-4-acetyloxy-4-(3-trifluoromethyl-4-chlorophenyl)piperidine.
- 30 5-chloro-1-[1-(6-methylaminocarbonylaminoindan-1-ylmethyl)piperidin-4-yl]-1*H*-indole

The acid addition salts of the invention are pharmaceutically acceptable salts of the compounds of Formula I formed with non-toxic acids. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, embonic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, 5 salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, and theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric acids.

10

The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of solutions for injection. For preparing such compositions, methods 15 well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients, or other additives normally used in the art may be used.

Conveniently, the compounds of the invention are administered in unit dosage form containing said compounds in an amount of about 0.01 to 100 mg.


20

The total daily dose is usually in the range of about 0.05 - 500 mg, and most preferably about 0.1 to 50 mg of the active compound of the invention.

The invention moreover relates to a method for the preparation of the novel 4-Aryl- 25 1-[amino(indan, dihydrobenzofuran or dihydrobenzothiophene)methyl]piperidines, -tetrahydropyridines or -piperazines of Formula I, comprising :

a) reacting an amino derivative of the following Formula II :

14



II

wherein R²-R⁸, X, Y, Z, Ar, and the dotted line are as previously defined, with a reagent of the formula R^{1'}-hal or R^{1'}-OCOR, in which formulas hal is halogen, R is alkyl, aryl or alkoxy and R^{1'} is acyl, thioacyl, a group R⁹VCO-, or a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- where R⁹, V, R¹⁰ and R¹¹ are as previously defined except that neither R¹⁰ nor R¹¹ may be hydrogen, or with a lower alkylsulfonyl halogenide, trifluoromethylsulfonyl halogenide or an isocyanate or thioisocyanate of the formula R¹⁰-N=C=O or R¹⁰-N=C=S wherein R¹⁰ is as previously defined;


10

- b) in order to prepare a compound of Formula I wherein R¹ is lower alk(en/yn)yl, cycloalk(en)yl, cycloalk(en)yl-lower alk(en/yn)yl or aryl-lower alkyl, alkylating an amino derivative of Formula II with an alkylating agent such as an alkylhalogenide R¹-hal, a mesylate R¹''OSO₂CH₃, a tosylate R¹''OSO₂C₆H₄-CH₃, or a similar alkylating reagent with suitable leaving groups, R¹ being lower alkyl, lower alkenyl, lower alkynyl, cycloalk(en)yl, cycloalk(en)yl-lower alk(en/yn)yl or aryl-lower alkyl;

15

- c) reducing the tetrahydropyridinyl double bond in derivatives of the following Formula III :

20




III

wherein R¹-R⁸, X, Y, and Ar are as previously defined; or

- d) alkylating an arylpiperazine, arylpiperidine, or aryltetrahydropyridine of the

15


formula V with an alkylating derivative of the formula IV :



5 wherein R¹-R⁸, X, Y, Z, Ar, and the dotted line are as previously defined, and w is a leaving group such as eg. halogen, mesylate, or tosylate; or

e) in order to obtain to form a compound of Formula I in which the substituents R¹ and R² together constitute a ring, ringclosure of a derivative of Formula VI :


10



in which R³-R⁸, X, Y, Z, Ar, m, Q, T and the dotted line are as previously defined and w is a leaving group such as halogen, mesylate, or tosylate;

15

f) in order to obtain a compound of Formula I in which R¹ is lower alk(en/yn)yl, cycloalk(en)yl, cycloalk(en)yl-lower alk(en/yn)yl or aryl-lower alkyl, reducing the carbonyl group of an amide derivative of the following Formula VII :




20

wherein $R^2 - R^8$, X, Y, Z, Ar and the dotted line are as previously defined and $R^{1''}$ is such a group that the group $R^{1''}CH_2$ constitutes a lower alk(en/yn)yl, cycloalk(en)yl, cycloalk(en)yl-lower alk(en/yn)yl or aryl-lower alkyl as embraced by the definition of R^1 ; or

5

g) introducing a substituent R^3 , R^4 or R^5 by reacting a compound of the following Formula VIII :




VIII

10 wherein one of $R^{3'}$ - R^5 is hydrogen and the other two are the corresponding R^3 , R^4 or R^5 as previously defined and R^1 , R^2 , R^6 - R^8 , X, Y, Z, Ar and the dotted line are as previously defined, by using a reactive reagent such as a halogen or a halogenating agent, a sulfonating agent, a nitration agent or a reactive agent generating carbonium ions (RCO^+ , R^+) wherein R is alkyl alkynyl, aryl cycloalkyl, or cycloalk(en/yn)yl; or

15

h) reducing the double bond in a compound of the following Formula IX :



IX


20

wherein $R^1 - R^8$, X, Y, Z, and Ar are as previously defined and one of the two dotted lines indicates a double bond; or

i) reducing the amide carbonyl in a compound of the following Formula X :

25

17



wherein R¹ - R⁵, R⁸, X, Y, Z, Ar and the dotted line are as previously defined.

- 5 whereupon the compound of Formula I is isolated as the free base or a pharmaceutically acceptable acid addition salt thereof.

The reaction in Method a) is conveniently performed at low temperature (eg. below room temperature) in an inert solvent such as acetone, dichloromethane, tetrahydro-
 10 furan or dimethoxyethane when reactive carboxylic acid chlorides, isocyanates, or isothiocyanates are used. Formylated amines are prepared from the corresponding amines by reaction in formic acid, with esters of formic acid, or by reaction with mixed formic acid anhydride prepared *in situ*. Generally reaction temperatures are between 0°C and the boiling point of the formyl precursor compounds.

15

The alkylations according to Methods b) and d) are generally performed by refluxing in a suitable solvent such as acetone, methyl isobutyl ketone, tetrahydrofuran, dioxane, ethanol or 2-propanol in the presence of a base such as triethylamine or potassium carbonate.

20

The reductions of double bonds according to Methods c) and h) are generally performed by catalytic hydrogenation at low pressure (< 3 atm.) in a Parr apparatus, or by using reducing agents such as diborane in inert solvents such as tetrahydrofuran, dioxane, or diethyl ether.

25

The reductions according to Methods f) and i) are generally performed by use of LiAlH₄, AlH₃ or diborane in an inert solvent such as tetrahydrofuran, dioxane, or diethyl ether at room temperature or at a slightly elevated temperature.

The halogenation according to Method g) is generally performed by use of chlorine, bromine, or N-chlorosuccinimide, N-bromosuccinimide or another halogen precursor molecule, conveniently in the presence of a catalyst such as Fe ions or a mineral acid.

5

1-Unsubstituted 4-arylpiperazines of Formula V ($Z = N$) are either commercially available or may be synthesized from the corresponding anilines and N',N'-bis(2-chloroethyl)amine by refluxing in high boiling solvents as chlorobenzene typically for 2-3 days according to methods described by Martin et al. *J.Med.Chem.* **1989**, *32*
10 1052-1056.

4-Arylpiperidines of formula V ($Z = CH$) are either commercially available or prepared as described in eg. US Pat. No. 2,891,066; McElvain et al. *J. Amer. Chem. Soc.* **1950**, *72*, 3134; Bally et al *Chem.Ber.* **1887**, *20*, 2590. The corresponding 4-aryl-1,2,3,6-tetrahydropyridines of Formula V ($Z = C$) are prepared from N-protected 4-piperidones by addition of properly substituted aryl lithium or magnesium halides followed by acid catalyzed water elimination. The N-protecting group (carbamate, benzyl, sulfonyl, acetyl) is finally removed in a conventional manner.

15

20 Synthesis of 3-(4-piperidinyl)-1*H*-indoles and 3-(1,2,3,6-tetrahydropyridin-4-yl)-1*H*-indoles is described in the Experimental Section.

Key intermediates such as 1-indancarboxylic acid (V. Asham and W.H. Linnell, *J. Chem. Soc.* **1954**, 4691-4693, Hansen et al. *Helv.Chim.Acta* **1982**, *33*, 325-343),
25 and 5-nitro-3-benzothiophenecarboxylic acid (EP Pat. Appln. No. 88-301073 CA(110(9):75302y (1988) and references cited therein) were prepared according to well-known literature procedures.

Experimental Section

30

In the following the invention is further illustrated by examples which, however, may not be construed as limiting.

In all the Examples, melting points were determined on a Büchi SMP-20 apparatus. Melting points are given as uncorrected values. ¹H NMR spectra were recorded at 250 MHz on a Bruker AC 250 spectrometer. Deuterated chloroform (99,8 %D) or dimethylsulfoxide (99,9 %D) were used as solvents. TMS was used as internal
5 reference standard. Chemical shift values are expressed in ppm-values. The following abbreviations are used for multiplicity of NMR signals: s=singlet, d=doub-
let, t=triplet, q=quartet, qui=quintet, h=heptet, dd=double doublet, dt=double triplet, dq=double quartet, tt=triplet of triplets, m=multiplet.

10 Example 1

6-Nitro-1-indancarboxylic acid, 1a

A solution of 1-indancarboxylic acid (30 g), prepared according to the method of Hansen et al. *Helv.Chim.Acta* **1982**, *33*, 325-343, in dichloromethane (50 ml) was
15 mixed with concentrated sulphuric acid (300 ml) at -10 °C. A mixture of 100 % HNO₃ (11.4 g) in concentrated H₂SO₄ (96 ml) was added dropwise under vigorous stirring below -10 °C. After stirring for one hour at 10 °C, the mixture was poured onto ice. Extraction with ethyl acetate (2x 300 ml), drying (anh. MgSO₄) and finally evaporation of the organic solvent afforded 42 g of the title compound. Mp : 126-
20 130 °C.

5-Nitro-3-benzothiophencarboxylic acid was prepared from 3-bromo-5-nitrobenzo-
thiophene via the corresponding 3-cyanobenzothiophene derivative according to EP
Pat. No. 88-301073 (CA110(9):75203y (1988), *J.Amer.Chem.Soc.* **1948**, *70*, 1955,
25 and *J.Chem.Soc.(c)* **1967**, 1899.

Example 2 (Method i)

1-(6-Aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, 2a

30 Dimethylformamide (DMF, 1 ml) was added to a solution of 6-nitro-1-indancarboxy-
lic acid, 1a (13 g) and thionylchloride (18 ml) in dichloromethane (125 ml). The mixture was heated to reflux for 4 hours. Toluene was added and volatile material was evaporated *in vacuo*. The thus obtained carboxylic acid chloride was dissolved

in dichloromethane (100 ml) and added dropwise to a solution of 4-(4-fluorophenyl)piperidine (19.5 g) and triethylamine (7 ml) in dichloromethane (100 ml) at 0-5 °C. The mixture was stirred at room temperature for another 1.5 hours. Water was added, the organic phase separated, washed with brine, dried (anh. MgSO₄),
5 filtered, and dichloromethane evaporated *in vacuo* leaving the crude 6-nitroindan-1-carboxamide derivative as an oil (35 g). Purification by column chromatography on silica gel (eluted with a 1:1 mixture of ethyl acetate and heptane) yielded 12 g of the pure carboxamide as an oil. All of this oil was dissolved in refluxing 90 % ethanol (350 ml). Fe-powder (10 g) and concentrated aqueous HCl (1 ml) were added
10 successively in small portions during 10 minutes. The resulting mixture was refluxed for another 2.5 hours. Inorganic salts were filtered off while still hot and ethanol evaporated *in vacuo*. Diluted aqueous NH₄OH was added until pH>9. Extraction with ethyl acetate (2x 200 ml) and working-up as above of the organic phase afforded 8 g of the 6-aminoindan-1-carboxamide derivative. Mp : 144-145 °C. To a
15 suspension of LiAlH₄ (2.7 g) in dry tetrahydrofuran (THF, 125 ml) was added dropwise a solution of all of the carboxamide in THF (125 ml). The mixture was gently refluxed for 2 hours. After cooling to 10 °C water (10 ml) and a 15% aqueous NaOH solution were cautiously added to destroy excess LiAlH₄. Inorganic salts were filtered off and washed extensively with THF. The combined THF solutions
20 were evaporated leaving 6.5 g of the title compound **2a** as an oil. The hydrochloride salt crystallized from 2-propanol. Mp : 198-201 °C. ¹H NMR (DMSO-d₆) : δ 1.85-2.40 (m, 6H); 2.60-2.90 (m, 3H); 3.00-3.15 (m, 3H); 3.35 (broad s, 3H); 3.45-3.60 (m, 2H); 3.65-3.75 (m, 1H); 6.45 (d, 1H); 6.50 (s, 1H); 6.95 (d, 1H); 7.15 (t, 2H); 7.25-7.35 (m, 2H).

25

In a similar manner the following aniline derivatives were prepared :

1-(6-Aminoindan-1-ylmethyl)-4-(2-methoxyphenyl)piperazine, **2b** as an oil

1-(6-Aminoindan-1-ylmethyl)-4-(2-chlorophenyl)piperazine, **2c** as an oil.

30 1-(6-Aminoindan-1-ylmethyl)-4-(3-chlorophenyl)piperazine, **2d** as an oil.

1-(6-Aminoindan-1-ylmethyl)-4-(4-chlorophenyl)piperazine, **2e**, mp : 98-109°C

1-(5-Amino-2,3-dihydrobenzothiophen-3-ylmethyl)-4-(4-fluorophenyl)piperidine, **2f** as an oil.

1-(6-Aminoindan-1-ylmethyl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine, **2g**, mp: 78-84 °C.

1-(6-Aminoindan-1-ylmethyl)-4-(3,4-dichlorophenyl)piperazine, **2h**, mp : 156-158 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.70-1.90 (m, 1H); 2.20-2.30 (m, 1H); 2.45 (dd, 1H); 2.55-2.75 (m, 6H); 2.75-2.90 (m, 2H); 3.20 (t, 4H); 3.20-3.35 (m, 1H); 3.55 (broad s, 2H); 6.50 (dd, 1H); 6.70-6.80 (m, 2H); 6.90-7.00 (m, 2H); 7.25 (d, 1H).

1-(6-Aminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperazine, **2i** as an oil
1-(6-Aminoindan-1-ylmethyl)-4-hydroxy-4-(3-trifluoromethyl-4-chlorophenyl)-
10 piperidine, **2j** as an oil

1-(6-Aminoindan-1-ylmethyl)-4-(4-methylphenyl)piperidine, **2k** as an oil.

1-(6-Aminoindan-1-ylmethyl)-4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridine, **2l** as an oil.

15 **Example 3 (Method i)**

1-(6-Aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperazine, **3a**

1-Indancarboxylic acid (20 g), DMF (2 ml), and thionylchloride (53 g) in dichloromethane (250) were refluxed for 4 hours. Volatile material was evaporated *in vacuo*
20 and remaining thionylchloride was removed by evaporation with toluene *in vacuo*. The remaining carboxylic acid chloride was dissolved in dichloromethane (200 ml) and added dropwise to a solution of 1-(4-fluorophenyl)piperazine (58 g) in dichloromethane (200 ml) at 0-5 °C. After stirring for 1.5 hours at room temperature, the organic phase was washed successively with water and brine and finally work-up as
25 above of the organic phase yielded 66 g crude carboxamide. Purification by column chromatography on silica gel (eluted with ethyl acetate / heptane 1:1) yielded 36 g of crystalline product with mp : 119-124 °C. All of this product was dissolved in concentrated H₂SO₄ (170 ml) at -10 °C. A mixture of 100 % HNO₃ (6.9 g) in concentrated H₂SO₄ (55 ml) was added dropwise under vigorous stirring below -10
30 °C. The mixture was stirred for another hour at -5 °C. The mixture was poured onto ice (500 g) and a 1:1 mixture of dichloromethane and ethyl acetate (300 ml) was added. The organic phase was separated and washed with diluted Na₂CO₃ solution (2 x 200 ml) and brine (200 ml). Work-up of the organic phase as above yielded 35

g of the crude 6-nitroindan-1-carboxamide derivative as an oil. The crude product was dissolved in 90% ethanol at reflux. Fe-powder (31.5 g) and concentrated aqueous HCl (3.1 ml) were added successively in small portions during 30 minutes. The resulting mixture was refluxed for another 2.5 hours. Inorganic salts were
5 filtered off while still hot and ethanol evaporated *in vacuo*. Diluted aqueous NH₄OH was added until pH>9. Extraction with dichloromethane (2x 200 ml) and working-up as above of the organic phase afforded 31 g of crystalline 6-aminoindan-1-carboxamide derivative. Mp : 143-149 °C. To a suspension of LiAlH₄ (10.4 g) in dry THF (400 ml) was added dropwise a solution of all of the carboxamide in THF (400 ml).
10 The mixture was gently refluxed for 2.5 hours. After cooling to 15 °C water (40 ml) and a 15% aqueous NaOH solution (10.4 ml) were cautiously added to destroy excess LiAlH₄. Inorganic salts were filtered off and washed extensively with THF. The combined THF solutions were evaporated leaving 23.7 g of the title compound **3a** as an oil.

15

Example 4 (method a)**1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine 4a**

To a solution of 1-(6-aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **2a** (6.5 g) and triethylamine (3 ml) in dichloromethane (150 ml) cooled to 0 °C was added
20 dropwise a solution of acetylchloride (1.7 g) in dichloromethane (50 ml). The mixture was stirred for one hour at room temperature. Water (500 ml) was added, the organic phase separated, washed with brine (2 x 50 ml) and finally worked-up as above. The thus isolated crude title product was purified by column chromatography on silica gel (eluted with a mixture of ethyl acetate/heptane/ triethylamine
25 75:25:4). Recrystallization from diethyl ether yielded 7.7 g of pure title compound **4a**. Mp : 159-162 °C. ¹H NMR (CDCl₃) : δ 1.70-1.90 (m, 5H); 2.00-2.15 (m, 1H); 2.15 (s, 3H); 2.20-2.30 (m, 1H); 2.35-2.50 (m, 2H); 2.60 (dd, 1H); 2.70-2.90 (m, 2H); 2.95-3.15 (m, 2H); 3.45 (qui, 1H); 6.95 (t, 2H); 7.05-7.25 (m, 5H); 7.55 (s, 1H)

30

In a corresponding manner, the following acylamino, thioacylamino, and sulfonylamino derivatives were prepared :

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperazine **4b**, mp : 179-187 °C (ethanol). ¹H NMR (DMSO-d₆) : δ 1.70-1.85 (m, 1H); 2.00 (s, 3H); 2.10-2.25 (m, 1H); 2.35 (dd, 1H), 2.50-2.60 (m, 5H); 2.65-2.90 (m, 2H); 3.10 (t, 4H); 3.35 (qui, 1H); 6.90-7.10 (m, 5H); 7.30 (d, 1H); 7.60 (s, 1H); 9.75 (s, 1H)

5

1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-4-(4-fluorophenyl)piperidine **4c**, mp : 140-142 °C (washed with diethyl ether). ¹H NMR (DMSO-d₆) : δ 1.55-1.75 (m, 4H); 1.95 (s, 3H); 1.90-2.05 (m, 1H); 2.05-2.20 (dt, 1H); 2.35 (dd, 1H); 2.40-2.55 (m, 3H); 2.95 (d, 1H), 3.15 (d, 1H); 3.20-3.35 (m, 1H); 3.45 (t, 1H); 3.55-10 3.70 (m, 1H); 7.05-7.15 (m, 3H); 7.25-7.35 (m, 3H); 7.65 (s, 1H); 9.85 (s, 1H)

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-chlorophenyl)piperazine **4d**, mp : 191-194 °C (acetone). ¹H NMR (CDCl₃) : δ 1.80-1.95 (m, 1H); 2.10 (s, 3H); 2.20-2.35 (m, 1H); 2.45 (dd, 1H); 2.60-2.70 (m, 5H); 2.80-2.95 (m, 2H); 3.15 (t, 4H); 3.35 (qui, 15 1H); 6.85 (d, 2H); 7.05-7.25 (m, 5H); 7.55 (s, 1H);

1-(6-Acetylaminoindan-1-ylmethyl)-4-(3-chlorophenyl)piperazine **4e**, mp : 176-178 °C (acetone). ¹H NMR (CDCl₃) : δ 1.75-1.90 (m, 1H), 2.15 (s, 3H); 2.20-2.35 (m, 1H); 2.45 (dd, 1H); 2.60-2.75 (m, 5H); 2.75-2.95 (m, 2H); 3.20 (t, 4H); 3.35 (qui, 1H); 20 6.80 (d, 2H); 6.85 (s, 1H); 7.10-7.30 (m, 4H); 7.55 (s, 1H)

1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-chlorophenyl)piperazine, hydrochloride, **4f** mp : 195-203 °C (acetone). ¹H NMR (DMSO-d₆) : δ 2.00 (s, 3H); 2.00-2.15 (m, 1H); 2.30-2.45 (m, 1H); 2.75-2.95 (m, 2H); 3.20-3.80 (m, 11H); 7.05-7.40 (m, 5H); 7.45 25 (d, 1H); 7.65 (s, 1H); 10.00 (s, 1H); 10.95 (broad s, 1H)

1-(6-Acetylaminoindan-1-ylmethyl)-4-(2-methoxyphenyl)piperazine, oxalate **4g**, mp : 210-213 °C (acetone/ethanol 1:1), ¹H NMR (DMSO-d₆) : δ 1.85-2.00 (m, 1H); 2.05 (s, 3H); 2.25-2.40 (m, 1H); 2.65-3.05 (m, 3H); 3.25 (broad s, 8H); 3.45-3.60 (m, 1H); 30 3.80 (s, 3H); 6.85-7.05 (m, 4H); 7.15 (d, 1H); 7.30 (d, 1H); 7.60 (s, 1H); 9.90 (s, 1H)

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridin **4h**,

mp : 156-161 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.80-1.95 (m, 1H); 2.15 (s, 3H); 2.20-2.35 (m, 1H); 2.45-2.60 (m, 3H); 2.65-3.00 (m, 5H); 3.20 (broad s, 2H); 3.30-3.45 (m, 1H); 6.05 (broad s, 1H); 6.95 (t, 2H); 7.15 (d, 1H); 7.15-7.25 (m, 2H); 7.35 (dd, 2H); 7.50 (s, 1H)

5

4-(4-Fluorophenyl)-1-(6-methansulfonylaminoindan-1-ylmethyl)piperidine **4i**, mp : 152-155 °C (diethyl ether), ¹H NMR (CDCl₃) : δ 1.70-1.90 (m, 5H); 2.00-2.20 (m, 2H); 2.20-2.35 (m, 1H); 2.40-2.70 (m, 3H); 2.75-2.95 (m, 2H); 3.00 (s, 3H); 3.10 (t, 2H); 3.25-3.45 (m, 1H); 6.70 (broad s, 1H); 6.90-7.05 (m, 3H); 7.15-7.25 (m, 3H);
10 7.35 (s, 1H)

1-(6-Cyclopropylcarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine **4j**, mp: 134-140 °C (diethyl ether), ¹H NMR (CDCl₃) : δ 0.75-0.90 (m, 2H); 1.05-1.15 (m, 2H); 1.45-1.60 (m, 1H); 1.75-1.95 (m, 6H); 2.00-2.15 (m, 2H); 2.20-2.35 (m, 1H);
15 2.35-2.50 (m, 2H); 2.65 (dd, 1H); 2.70-2.95 (m, 2H); 3.00-3.15 (m, 2H); 3.35 (qui, 1H); 6.95 (t, 2H); 7.05-7.25 (m, 4H); 7.40 (broad s, 1H); 7.65 (broad s, 1H)

1-(6-Cyclopentylcarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine **4k**, mp : 177-178 °C (diethyl ether). ¹H NMR (CDCl₃) : δ 1.50-1.70 (m, 2H); 1.70-1.95
20 (m, 11H); 2.00-2.15 (m, 2H); 2.20-2.35 (m, 1H); 2.35-2.50 (m, 2H); 2.60-2.75 (m, 2H); 2.75-2.95 (m, 2H); 2.95-3.15 (m, 2H); 3.35 (qui, 1H); 6.95 (t, 2H); 7.10-7.25 (m, 5H); 7.65 (s, 1H)

1-(6-Ethylloxycarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, fumarate **4l**, mp: 191-193 °C (ethanol/acetone 2:1). ¹H NMR (DMSO-d₆) : δ 1.15 (s, 3H); 1.70-
25 1.90 (m, 5H); 2.10-2.30 (m, 3H); 2.40-2.90 (m, 5H); 3.10-3.20 (m, 2H); 3.25-3.35 (m, 1H); 4.15 (q, 1H); 6.60 (s, 1.5H); 7.00-7.15 (m, 4H); 7.35 (dd, 2H); 7.55 (s, 1H); 9.45 (s, 1H).

4-(4-Chlorophenyl)-1-(6-methansulphonylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridine, oxalate **4m**, mp : 176-178 °C (from ethanol), ¹H NMR
30 (DMSO-d₆) : δ 1.90-2.00 (m, 1H); 2.30-2.40 (m, 1H); 2.95 (s, 3H); 2.65-3.10 (m, 5H); 3.20-3.30 (m, 3H); 3.50-3.60 (m, 1H); 3.75 (broad s, 2H); 6.25 (broad s, 1H); 7.05 (dd, 1H); 7.15-7.25 (m, 2H); 7.45 (d, 2H); 7.55 (d, 2H); 9.60 (broad s, 1H).

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-methylphenyl)piperidine **4n**, mp : 173-175 °C (washed with diethyl ether), ¹H NMR (CDCl₃) : δ 1.70-1.95 (m, 5H); 2.10 (s, 3H); 2.20-2.30 (m, 1H); 2.30 (s, 3H); 2.35-2.50 (m, 2H); 2.65 (dd, 1H); 2.70-2.90 (m, 2H); 3.00-3.15 (m, 2H); 3.35 (qui, 1H); 7.05-7.25 (m, 6H); 7.50 (s, 1H); 7.55 (s, 1H).

1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-dichlorophenyl)piperazine **4o**, mp : 160-163 °C (washed with diethyl ether), ¹H NMR (CDCl₃) : δ 1.80-1.90 (m, 1H); 2.10 (s, 3H); 2.15-2.30 (m, 1H); 2.45 (dd, 1H); 2.55-2.70 (m, 5H); 2.70-3.00 (m, 2H); 3.20 (t, 4H); 3.35 (qui, 1H); 6.25 (dd, 1H); 6.95 (d, 1H); 7.10 (d, 1H); 7.20 (dd, 1H); 7.25 (d, 1H); 7.40 (broad s, 1H); 7.60 (s, 1H).

1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridine, oxalate **4p**, mp : 223-226 °C (from acetone), ¹H NMR (DMSO-d₆) : δ 1.80-1.95 (m, 1H); 2.05 (s, 3H); 2.20-2.40 (m, 1H); 2.65-3.00 (m, 5H); 3.15-3.30 (m, 3H); 3.50-3.60 (m, 1H); 3.70 (broad s, 2H); 6.25 (broad s, 1H); 7.15 (d, 1H); 7.30 (d, 1H); 7.40 (d, 2H); 7.50 (d, 2H); 7.65 (s, 1H).

1-(6-Acetylaminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperazine, **4q**, mp : 188-189 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.70-1.95 (m, 1H); 2.15 (s, 3H); 2.15-2.30 (m, 1H); 2.45 (dd, 1H); 2.60-2.70 (m, 5H); 2.70-2.90 (m, 2H); 3.10 (t, 4H); 3.40 quin, 1H); 5.85 (s, 2H); 6.45 (dd, 1H); 6.55 (d, 1H); 6.70 (d, 1H); 7.15 (d, 1H); 7.20-7.35 (m, 2H); 7.55 (s, 1H).

1-(6-Acetylaminoindan-1-ylmethyl)-4-hydroxy-4-(3-trifluoromethyl-4-chlorophenyl)piperidine, hemioxalate, **4r**, mp : 163-165 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.65-1.95 (m, 3H); 2.05 (s, 3H); 2.15-2.30 (m, 3H); 2.65-3.30 (m, 8H); 3.40-3.50 (m, 1H); 7.15 (d, 1H); 7.25 (d, 1H); 7.70-7.85 (m, 3H); 8.00 (s, 1H).

Example 5 (method a)

4-(4-Fluorophenyl)-1-(6-methylaminocarbonylaminoindan-1-ylmethyl)piperidine, **5a**

1-(6-Aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **2a** (3 g) was dissolved in dichloromethane (140 ml) and methylisocyanate (0.53 g) was added. The solution was refluxed for 4 hours. Dichloromethane was evaporated and the remaining

crude title compound was purified by column chromatography on silica gel (eluted with 4% triethylamine in ethyl acetate). The pure title compound **5a** crystallized from diethyl ether. Yield : 0.8 g, mp : 170-173 °C. ¹H NMR (CDCl₃) : δ 1.70-1.90 (m, 5H); 2.00-2.15 (m, 2H); 2.20-2.35 (m, 1H); 2.35-2.55 (m, 2H); 2.60 (dd, 1H); 2.80 (d, 3H);
 5 2.75-2.95 (m, 2H); 3.05 (broad d; 2H); 3.30 (qui, 1H); 4.95 (q, 1H); 6.55 (s, 1H); 6.90-7.00 (m, 3H); 7.10-7.25 (m, 3H); 7.35 (s, 1H)

In a similar manner the following urea or thiourea derivatives were prepared :

1-[6-(4-Fluorophenyl)aminocarbonylaminoindan-1-ylmethyl]-4-(4-fluorophenyl)-
 10 piperidine, **5b**, mp : 235-238 °C (CH₂Cl₂). ¹H NMR (DMSO-d₆) : δ, 1.55-1.80 (m, 5H); 2.00-2.30 (m, 3H); 2.35 (dd, 1H); 2.45-2.60 (m, 2H); 2.60-2.85 (m, 2H); 3.05 (broad d, 2H); 3.30 (qui, 1H); 7.05-7.15 (m, 6H); 7.25-7.35 (dd, 2H); 7.40-7.50 (dd, 2H); 7.55 (s, 1H); 8.50 (s, 1H); 8.65 (s, 1H)

4-(4-Fluorophenyl)-1-(6-methylaminothiocarbonylaminoindan-1-ylmethyl)piperidine,
 15 fumarate, **5c**, mp: 181-183°C (ethanol/acetone 1:1). ¹H NMR (DMSO-d₆) : δ 1.70-1.90 (m, 5H); 2.15-2.95 (m, 8H); 2.90 d, 3H); 3.20-3.30 (m, 2H); 3.40 (qui, 1H); 6.20 (s, 2H); 7.05-7.20 (m, 4H); 7.30 (dd, 2H); 7.40 (s, 1H); 7.80 (broad s, 1H); 9.60 (broad s, 1H).

1-(6-Methylaminocarbonylaminoindan-1-ylmethyl)-4-(3,4-methylenedioxyphenyl)piperazine, hemioxalate, **5d**, mp : 132-133 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.70-1.85 (m, 1H); 2.15-2.30 (m, 1H); 2.60 (d, 3H); 2.70-3.00 (m, 7H); 3.20 (m, 4H); 3.35-3.45 (m, 1H); 5.90 (s, 2H); 6.00-6.10 (m, 1H); 6.40 (dd, 1H); 6.70 (d, 1H); 6.80 (d, 1H); 7.05 (d, 1H); 7.10 (d, 1H); 7.45 (s, 1H); 8.40 (s, 1H).

25 **Example 6 (method a)**

1-(6-Dimethylaminocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **6a**

1-(6-Aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **2a** (3 g) was dissolved in THF (50 ml) and triethylamine (2 g) was added. At 5 °C dimethylcarbonylchloride
 30 (1 g) in THF (15 ml) was added dropwise. After completed addition the mixture was refluxed for 1.5 hours. THF was evaporated. Water was added and extraction with dichloromethane (2 x 50 ml) and work-up as above of the combined organic phases

yielded crude title product, which was purified by column chromatography on silica gel (eluted with 4% triethylamine in a 1:1 mixture of ethyl acetate and heptane). The pure title compound **6a** crystallized from diethyl ether. Yield : 1.4 g, mp : 141-144 °C. ¹H NMR (CDCl₃) : δ 1.65-2.55 (m, 10H); 2.70 (dd, 1H); 2.70-2.95 (m, 2H); 2.95-3.05 (m, 1H); 3.05 (s, 6H); 3.15 (broad d, 1H); 3.35 (qui, 1H); 6.25 (s, 1H); 6.95-7.25 (m, 6H); 7.45 (s, 1H)

In a similar manner the following urea and thiourea derivatives were prepared :

1-(6-Dimethylaminothiocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine
10 **6b**, mp : 175-180 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.65-1.95 (m, 5H); 2.00-2.15 (m, 2H); 2.20-2.35 (m, 1H); 2.35-2.55 (m, 2H); 2.60 (dd, 1H); 2.70-2.90 (m, 2H); 3.05-3.15 (m, 2H); 3.85 (s, 6H); 3.40 (qui, 1H); 6.95-7.10 (m, 4H); 7.15-7.30 (m, 4H)

4-(4-fluorophenyl)-1-[6-(1-pyrrolidinyl)carbonylaminoindan-1-ylmethyl]piperidine **6c**,
15 mp: 190-193 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.65-2.55 (m, 14H); 2.65 (dd, 1H); 2.75-2.90 (m, 2H); 3.00 (broad d, 1H); 3.15 (broad d, 1H); 3.35 (qui, 1H); 3.45 (t, 4H); 6.15 (s, 1H); 6.95 (t, 2H); 7.00-7.10 (m, 2H); 7.20 (dd, 2H); 7.50 (s, 1H)

20 **Example 7** (method a)

1-(6-Aminocarbonylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, hemifumarate, **7a**

A solution of potassium isocyanate (1.5 g) dissolved in dichloromethane (20 ml)
25 was cooled to 5 °C and a solution of trifluoroacetic acid (1.9 g) in dichloromethane (20 ml) was added dropwise. To the resulting mixture was added dropwise a solution of 1-(6-aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **2a** (3 g) in dichloromethane (10 ml). The temperature was allowed to raise to room temperature. After stirring for another 3 hours the mixture was poured on ice (500 g) and
30 diluted aqueous NH₄OH was added until pH > 9. The organic phase was separated and worked-up as above. The crude title product was purified by column chromatography on silica gel (eluted with 4% triethylamine in ethanol / ethyl acetate 1:3). The

purified product (2 g) was dissolved in acetone (20 ml) and added to a solution of fumaric acid (0.6 g) in ethanol (20 ml). The precipitated hemifumarate salt was filtered off and dried. Yield : 1.6 g, mp: 172-174 °C. ¹H NMR (DMSO-d₆) : δ 1.65-1.90 (m, 5H); 2.10-2.35 (m, 3H); 2.45-2.90 (m, 5H); 3.10-3.25 (m, 2H); 3.45 (qui,
5 1H); 6.85 (s, 2H); 6.60 (s, 1H); 7.00-7.15 (m, 4H); 7.30 (dd, 2H); 7.45 (s, 1H); 8.45 (s, 1H).

Example 8

5-Chloro-3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indole, 8a

10

A mixture of 5-chloro-1H-indole (25 g), piperidin-4-one,hydrate,hydrochloride (71 g), and potassium hydroxide (38 g) in ethanol (450 ml) was refluxed for 6 hours. After cooling inorganic salts were filtered off and ethanol evaporated *in vacuo*. To the remaining oil was added brine (500 ml) and ethyl acetate (2 x 200 ml). The organic
15 phase was separated and worked-up as above. Yield of crude title product : 45 g (semicrystalline).

In a corresponding manner the following 3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indoles were prepared :

20 6-Chloro-3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indole, 8b

5-Fluoro-3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indole, 8c

Example 9

5-Chloro-3-(4-piperidiny)-1H-indole, 9a

25

Crude 5-chloro-3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indole 8a (26 g) was dissolved in glacial acetic acid (330 ml) and PtO₂ (0.7 g) was added. The mixture was hydrogenated in a Parr apparatus at 3 atm. for 5 hours. The catalyst was filtered off and excess acetic acid evaporated *in vacuo*. Water was added and pH was
30 adjusted to >9 by addition of diluted aqueous NH₄OH. Extraction with ethyl acetate (2 x 200 ml) and work-up of the combined organic phases yielded 19 g of crude title compound as a visceous oil.

In a corresponding manner the following 3-(4-piperidiny)-1 *H*-indoles were prepared:

6-Chloro-3-(4-piperidiny)-1 *H*-indole, **9b**

5-Fluoro-3-(4-piperidiny)-1 *H*-indole, **9c**

5 Example 10

3-[1-(5-Amino-2,3-dihydrobenzothiophen-3-ylmethyl)piperidin-4-yl]-5-chloro-1 *H*-indole, **10a**

5-Nitro-3-benzothiophencarboxylic acid **1b** (20 g) was converted into the corresponding carboxylic acid chloride as in Example 2. The acid chloride was dissolved in THF (200 ml) and added dropwise to a solution of 5-chloro-3-(4-piperidiny)-1 *H*-indole, **9a** (19 g) and triethylamine (10 ml) in THF (200 ml) at 0-5 °C. The mixture was stirred overnight at room temperature. THF was evaporated. Water was added to the remaining oil. Extraction with dichloromethane (2 x 100 ml) and work-up of the organic extracts afforded the crude 5-nitro-3-benzothiophencarboxylic acid amide, which was subsequently purified by column chromatography on silica gel (eluted with ethyl acetate/heptane 1:1). Yield 6.6 g, mp : 243-250 °C. All of the amide was dissolved in 90% ethanol at reflux. Fe-powder (5 g) and concentrated aqueous HCl were added successively in small portions during 10 minutes. The mixture was refluxed for another 2.5 hours. Inorganic salts were filtered off and ethanol evaporated *in vacuo*. Water was added to the remaining oil and pH was adjusted to >9 by addition of diluted aqueous NH₄OH. Extraction with dichloromethane (2 x 100 ml) and subsequent work-up of the organic phase yielded 4 g of the 5-amino-3-benzothiophencarboxylic acid amide as an oil. All of this oil was dissolved in methanol (100 ml). 0.5 g of Mg turnings were added. By heating to 35 °C an exothermic reaction started. Mg turnings were added in small portions (3x0.5g) while keeping the temperature below 45 °C. The mixture was finally poured into an aqueous NH₄Cl solution and concentrated aqueous HCl (1 ml) was added. Extraction with dichloromethane (2 x 50 ml) and work-up of the organic extracts as above afforded 2 g of the 5-amino-2,3-dihydrobenzothiophen-3-carboxylic acid amide as an oil. To a suspension of LiAlH₄ (0.6 g) in dry THF (50 ml) was added dropwise a solution of all of the carboxamide in THF (50 ml). The mixture was gently refluxed for 2 hours. After cooling to 10 °C water (2.4 ml) and a

15% aqueous NaOH solution were cautiously added to destroy excess LiAlH₄. Inorganic salts were filtered off and washed extensively with THF. The combined THF solutions were evaporated leaving 1.9 g of 3-[1-(5-amino-2,3-dihydrobenzothiophen-3-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indole as an oil.

5

In a similar manner the following aniline derivatives were prepared :

3-[1-(5-Amino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1*H*-indole, **10b** as an oil

3-[1-(5-Amino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-
10 5-fluoro-1*H*-indole, **10c** as an oil.

The corresponding 1-(6-Aminoindan-1-ylmethyl)-substituted 4-(3-indolyl)piperidines and 4-(3-indolyl)-1,2,3,6-tetrahydropyridines were prepared from the corresponding 1-indancarboxamides, which were successively nitrated in the 6 position, reduction
15 of the nitro substituent and reduction of the carboxamide carbonyl group. A reaction sequence as outlined in Example 3. The following indan derivatives were prepared accordingly :

3-[1-(6-Aminoindan-1-ylmethyl)piperidin-4-yl]-5-fluoro-1*H*-indole, **10d** as an oil.

3-[1-(6-Aminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1*H*-indole, **10e** as an oil.

20 3-[1-(6-Aminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1*H*-indole, **10f** as an oil.

3-[1-(6-Aminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1*H*-indole, **10g** as an oil.

1-[1-(6-aminoindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indole, **10h** as an oil.

25

Example 11

3-[1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indole, oxalate, **11a**

30 To a solution of 3-[1-(5-amino-2,3-dihydrobenzothiophen-3-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indole **10a** (1.9 g) and triethylamine (2 ml) in dichloromethane (50 ml) kept at 0 °C was added dropwise a solution of acetylchloride (0.4 g) in dichloromethane (10 ml). The mixture was stirred at room temperature for 2 hours. Water was

added and the organic phase was worked-up as above. The crude title compound was purified by column chromatography on silica gel (luted with 4% triethylamine in ethyl acetate). Yield : 0.8 g. The oxalate salt of the title compound crystallized from a 1:1 mixture of acetone and ethanol. Mp: 168-174 °C. ¹H NMR (DMSO-d₆) : δ

5 2.00 (s, 3H); 1.95-2.15 (m, 4H); 2.85-3.25 (m, 5H); 3.40-3.50 (m, 2H); 3.55-3.70 (m, 2H); 3.90-4.00 (m, 1H); 7.00-7.40 (m, 6H); 7.70 (s, 1H); 7.70 (broad s, 2H); 9.95 (broad s, 1H); 11.10 (broad s, 1H).

In a similar manner the following indolyl derivatives were prepared :

- 10 3-[1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1*H*-indole, oxalate **11b**, mp : 214-216 °C (ethanol), ¹H NMR (DMSO-d₆) : δ 2.00 (s, 3H); 2.75 (broad s, 2H); 2.95-3.40 (m, 5H); 3.50-3.80 (m, 3H); 3.95 (broad s, 1H); 6.15 (broad s, 1H); 7.15 (t, 2H); 7.25 (d, 1H); 7.45 (d, 1H); 7.60 (s, 1H); 7.65 (s, 1H); 7.85 (s, 1H); 9.95 (s, 1H); 11.50 (s, 1H)
- 15 3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-5-fluoro-1*H*-indole, oxalate **11c**, mp : 145-149 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.80-2.15 (m, 6H); 2.00 (s, 3H); 2.30-2.45 (m, 1H); 2.70-2.90 (m, 2H); 2.95-3.20 (m, 3H); 3.35 (d, 1H); 3.50-3.80 (m, 3H); 6.95 (dt, 1H); 7.15 (d, 1H); 7.25 (s, 1H); 7.30 (d, 1H); 7.40 (dd, 1H); 7.45 (dd, 1H); 7.70 (s, 1H); 9.95 (s, 1H); 11.05 (s, 1H)
- 20 3-[1-(5-Acetylamino-2,3-dihydrobenzothiophen-3-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-fluoro-1*H*-indole, oxalate **11d**, mp : 155-165 °C (acetone). ¹H NMR (DMSO-d₆) : δ 2.00 (s, 3H); 2.75 (broad s, 2H); 2.95-3.45 (m, 5H); 3.50-3.80 (m, 3H); 3.95 (broad s, 1H); 6.15 (broad s, 1H); 6.95 (t, 1H); 7.20 (d, 1H); 7.30 (d, 1H); 7.45 (m, 1H); 7.55-7.70 (m, 3H); 9.95 (s, 1H); 11.45 (s, 1H)
- 25 3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1*H*-indole, oxalate hemihydrate **11e**, mp : 151-164 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.95-2.10 (m, 1H); 2.00 (s, 3H); 2.30-2.45 (m, 1H); 2.70-2.90 (m, 4H); 3.15 (t, 1H); 3.35-3.50 (m, 3H); 3.55-3.70 (m, 1H); 3.95 (broad s, 2H); 6.15 (s, 1H); 7.05 (dd 1H); 7.15 (d, 1H); 7.25 (d, 1H); 7.45 (d, 1H); 7.55 (d, 1H); 7.60 (s, 1H); 7.85 (d, 1H); 9.95
- 30 (s, 1H); 11.55 (s, 1H)
- 3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1*H*-indole, oxalate **11f**, mp : 122-130 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.90-2.15 (m, 6H); 2.00 (s, 3H);

2.25-2.40 (m, 1H); 2.70-3.10 (m, 6H); 3.35 (d, 1H); 3.45-3.65 (m, 2H); 7.00 (dd, 1H); 7.15-7.25 (m, 2H); 7.30 (d, 1H); 7.40 (s, 1H); 7.60-7.70 (m, 2H); 9.90 (s, 1H); 11.05 (s, 1H)

3-[1-(6-Acetylaminindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-5-chloro-1*H*-

5 indole, oxalate **11g**, mp : 220-223 °C (acetone/ethanol 5:1). ¹H NMR (DMSO-*d*₆) : δ 1.95-2.10 (m, 1H); 2.00 (s, 3H); 2.30-2.45 (m, 1H); 2.70-2.95 (m, 4H); 3.10 (t, 1H); 3.30-3.45 (m, 3H); 3.55-3.70 (m, 1H); 3.85 (broad s, 2H); 6.15 (s, 1H); 7.10-7.20 (m, 2H); 7.30 (d, 1H); 7.45 (d, 1H); 7.60-7.70 (m, 2H); 7.85 (s, 1H); 9.90 (s, 1H); 11.50 (s, 1H)

10 1-[1-(6-Acetylaminindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1*H*-indole, **11h**, mp: 189-191 °C (ethyl acetate). ¹H NMR (CDCl₃) : δ 1.80-2.00 (m, 1H); 2.05-2.40 (m, 7H); 2.20 (s, 3H); 2.50 (dd, 1H); 2.65 (dd, 1H); 2.80-2.95 (m, 2H); 3.15 (broad t, 2H); 3.35 (quin, 1H); 4.20-4.30 (m, 1H); 6.45 (d, 1H); 7.20-7.35 (m, 4H); 7.30-7.40 (m, 2H); 7.60 (d, 1H); 7.70 (broad s, 1H).

15

Example 12 (method e)

4-(4-Fluorophenyl)-1-[6-(1-pyrrolidin-2-onyl)indan-1-ylmethyl]piperidine, fumarate, **12a**

20 To a solution of 1-(6-aminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine, **2a** (3 g) and triethylamine (2 ml) in dichloromethane (50 ml) at 0 °C was added dropwise a solution of 4-chlorobutyric acid chloride (1.4 g) in dichloromethane (15 ml). The mixture was finally stirred for 5 hours at room temperature. Ice cold diluted aqueous NaOH solution was added and the organic phase was subsequently worked-up as
25 above. The crude 1-[6-(4-chlorobutanoylamino)indan-1-ylmethyl]-4-(4-fluorophenyl)-piperidine was purified by column chromatography on silica gel (eluted with 4% triethylamine in a 1:3 mixture of ethyl acetate and heptane). Yield of crystalline product : 2.4 g with mp: 129-135 (washed with diethyl ether). A solution of the thus isolated 4-chlorobutanoyl derivative (1 g) and potassium *tert*-butoxide (0.4 g) in dry
30 THF (40 ml) was refluxed for 2 hours. THF was evaporated *in vacuo*. Diluted aqueous NH₄OH and dichloromethane were added and the organic phase was subsequently worked-up as above. The remaining oil (1 g) was dissolved in acetone (10 ml) and added to a hot solution of fumaric acid (0.3 g) in ethanol (15 ml). After

cooling in a refrigerator overnight the precipitated fumarate salt was filtered off and dried. Yield : 0.7 g, mp: 177-179 °C. ¹H NMR (DMSO-d₆) : δ 1.70-1.90 (m, 5H); 2.10 (qui, 2H); 2.20-2.40 (m, 3H); 2.45-2.65 (m, 4H); 2.70-2.95 (m, 3H); 3.20 (broad t, 2H); 3.40 (qui, 1H); 3.70-3.90 (m, 2H); 6.60 (s, 2H); 7.15 (t, 2H); 7.20 (d, 1H); 7.30
5 (dd, 2H); 7.40 (dd, 1H); 7.65 (s, 1H).

EXAMPLE 13

(+)-6-Nitro-1-indancarboxylic acid, **13a**

10 A solution of 6-nitro-1-indancarboxylic acid (**1a**) (96 g) and brucine dihydrate (200 g) was heated in acetone (1.25 L) until a clear solution was obtained. The solution was left in a refrigerator overnight. The precipitated brucine salt was filtered off. Yield : 159.1 g. Recrystallization from 2-propanol afforded 103 g of pure (+)-6-nitro-1-indancarboxylic acid brucine salt. The salt was dissolved in water and diluted
15 hydrochloric acid was added. Extraction with diethyl ether and work-up as above afforded 29.8 g of **13a**. Mp : 92-94 °C. [α]_D = +83.3 ° (c = 1, methanol).

EXAMPLE 14

(+)-1-(6-Acetylaminoindan-1-ylmethyl)-4-(4-fluorophenyl)piperidine **14a**

20

The (+)-enantiomer of compound **4a** was prepared from (+)-6-Nitro-1-indancarboxylic acid, **13a** according to the methods in Examples 2 and 4. Mp : 145-146 °C. ¹H NMR (CDCl₃) : δ 1.70-1.90 (m, 5H); 2.00-2.15 (m, 1H); 2.15 (s, 3H); 2.20-2.30 (m, 1H); 2.35-2.50 (m, 2H); 2.60 (dd, 1H); 2.70-2.90 (m, 2H); 2.95-3.15 (m, 2H); 3.45
25 (qui, 1H); 6.95 (t, 2H); 7.05-7.25 (m, 5H); 7.55 (s, 1H). [α]_D = +24.3 ° (c = 1, methanol).

EXAMPLE 15

3-[1-(6-Acetylaminoindan-1-ylmethyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1-methyl-1*H*-indole **15a**
30

3-[1-(6-nitroindan-1-ylkarbonyl)-1,2,3,6-tetrahydropyridin-4-yl]-6-chloro-1*H*-indole

(23 g), prepared according to the procedure in Example 10, was dissolved in dry DMF (300 ml) and potassium tert-butoxide (7.3 g) was added at 10 °C. Methyl iodide (23.2 g) was added dropwise during 30 min. The mixture was left at room temperature overnight. Water and diethyl ether were added and the organic phase
5 was worked-up as above. The crude N-methylated indole was purified by column chromatography on silica gel (eluted with a 1:1 mixture of ethyl acetate and heptane). Yield 2.75 g. Reduction of the nitro group with Fe in 90 % acidic ethanol, followed by reduction of the amide carbonyl group and finally acetylation of the anilino group according to the methods in Examples 10 and 11 affords the title
10 compound **15a**. Mp : 189-193 °C (washed with diethyl ether). ¹H NMR (CDCl₃) : δ 1.80 (broad s, 1H); 1.80-1.95 (m, 1H); 2.10 (s, 3H); 2.25-2.40 (m, 1H); 2.50-2.60 (m, 3H); 2.70-2.95 (m, 5H); 3.25 (broad s, 2H); 3.40 (qui, 1H); 3.70 (s, 3H); 6.15 (broad s, 1H); 7.00 (s, 1H); 7.10 (dd, 1H); 7.15 (d, 1H); 7.20-7.30 (m, 3H); 7.50 (s, 1H); 7.80 (d, 1H).

15

Catalytic hydrogenation of compound **15a** according to the method in Example 9 afforded :

3-[1-(6-Acetylaminoindan-1-ylmethyl)piperidin-4-yl]-6-chloro-1-methyl-1*H*-indole, oxalate **15b**, mp : 202-205 °C (acetone). ¹H NMR (DMSO-d₆) : δ 1.90-2.15 (m, 5H);
20 2.00 (s, 3H); 2.30-2.40 (m, 1H); 2.70-3.15 (m, 6H); 3.35 (d, 1H); 3.50-3.70 (m, 2H); 3.75 (s, 3H); 7.00 (dd, 1H); 7.10-7.30 (m, 3H); 7.55 (d, 1H); 7.65 (s, 1H); 7.65 (d, 1H); 9.90 (s, 1H).

EXAMPLE 16 (Method f)

25 4-(4-Fluorophenyl)-1-(6-methylaminoindan-1-ylmethyl)piperidine, 1.5 oxalate **16a**

To a solution of 4-(4-fluorophenyl)-1-(6-methylaminoindan-1-ylmethyl)piperidine, **2a** (4 g) and triethylamine (3 mL) in dichloromethane was added dropwise at 0-5 °C a solution of ethyl chloroformate (1.5 g) in dichloromethane (15 mL). The mixture was
30 stirred at room temperature for 2 hours and poured onto saturated brine (500 mL). The organic phase was separated and worked-up as previously. Yield of the ethyl carbamate as an oil 4.3 g. To a suspension of LiAlH₄ (1.2 g) in dry diethyl ether (20 mL) at 5 °C was added dropwise a solution of all of the ethyl carbamate in dry THF

(25 mL). The mixture was stirred for an additional hour at 5 °C and finally at room temperature for 5 hours. Excess of LiAlH_4 was destroyed by cautiously adding water and diluted aqueous NaOH solution (6 mL). Precipitated inorganic salts were filtered off and the solvents evaporated *in vacuo*. The crude title compound was
5 purified by column chromatography on silica gel (eluted with a 1:1 mixture of heptane and ethyl acetate). Yield as an oil 1.5 g. The 1.5 oxalate salt **16a** crystallized from a 1:1 mixture of acetone and ethanol. Mp : 84-86 °C. ^1H NMR (DMSO- d_6) : δ 1.75-2.10 (m, 5H); 2.20-2.40 (m, 1H); 2.60 (s, 3H); 2.60-2.90 (m, 3H); 3.00-3.15 (m, 3H); 3.40-3.80 (m, 4H); 6.40(dd, 1H); 6.50 (d, 1H); 6.95 (d, 1H);
10 7.15 (t, 2H); 7.35 (dd, 2H)

EXAMPLE 17

5-Chloro-1-(4-piperidiny)-1*H*-indole, **17a**

15 To a solution of 5-chloro-1*H*-indole (20 g) in N-methyl-2-pyrrolidone (450 mL) were added potassium carbonate (82 g), CuBr (7.5 g), and Cu bronze (3 g). The mixture was heated to 140 °C and 4-bromopyridine, hydrochloride (22 g). The mixture was heated for 1 hour at 150 °C and further 4-bromopyridine, hydrochloride (15 g) was added. This procedure was repeated twice and the mixture was finally heated
20 overnight at 150 °C. After cooling precipitated inorganic salts were filtered off. Water (2 L), ethyl acetate (500 mL), and diluted aqueous ammonia (200 mL) were added. Undissolved material was filtered off and discarded. The organic phase was worked-up as above affording 34 g of 5-chloro-1-(4-pyridyl)-1*H*-indole with mp : 153-155 °C. All of this product, without further purification, was dissolved in
25 dimethoxyethane (350 mL) at 60-70 °C. Methyl iodide (14 mL) was added and the mixture was heated at reflux for 7 hours. After cooling the precipitated quaternized pyridinium salt was filtered off and washed with dimethoxyethane. Yield 32 g. Mp : 257-260 °C. All of the pyridinium salt was susoended in ethanol (450 mL) and water (50 mL). NaBH_4 (16 g) was added in small portions during 1.5 hours with stirring.
30 After stirring for another 1.5 hours most of the ethanol was evaporated at room temperature *in vacuo*. Ethyl acetate (300 mL) and water (500 mL) were added and the organic was worked up as previously. Yield of 5-chloro-1-(1-methyl-1,2,3,6-tetrahydro-4-pyridinyl)-1*H*-indole, 17 g as an oil. To a solution of the

tetrahydropyridinyl (15 g) derivativ in glacial acetic acid (150 mL) was added PtO_2 and the mixture was hydrogenated in a Parr apparatus at 3 ato for 7 hours. The catalyst was filtered off, most of the acetic acid evaporated *in vacuo* and finally the crude 5-chloro-1-(1-methyl-4-piperidinyl)-1*H*-indole was extracted with ethyl acetate
5 from an alkaline aqueous solution. Yield 13 g as an oil. Any remaining water in this crude product (10 g) was removed by evaporation with toluene. Finally the oil was dissolved in 1,1,1-trichloroethane (200 mL). At reflux temperature 2,2,2-trichloroethyl chloroformate (6.5 mL) dissolved in 1,1,1-trichloroethane (20 mL) was added dropwise. The mixture was refluxed for another 2 hours, sodium carbonate
10 (2 g) was added and reflux continued for 0.5 hours. After cooling the mixture was filtered through silicagel (eluted with dichloromethane). Yield 10 g of crude carbamate derivative. To a solution of the carbamate (6 g) in 90 % aqueous acetic acid (110 mL) was added finely powdered Zn (12 g) in small portions at 45 °C during 1 hour. The mixture was heated for another hour at 50 °C, Zn-salts were
15 filtered off and most of the acetic acid evaporated *in vacuo*. The remaining oil was dissolved in water (200 mL) and ethyl acetate (200 mL). The pH of the aqueous phase was adjusted >9 by adding diluted aqueous ammonia. The organic phase was finally worked-up as previously yielding 3.5 g of crude title product 17a as an oil which was used without further purification to prepare compound 10h.

20

EXAMPLE 18

1-(6-Acetylaminoindan-1-ylmethyl)-4-acetyloxy-4-(3-trifluoromethyl-4-chlorophenyl)piperidine, 18a.

25 To a solution of 1-(6-aminoindan-1-ylmethyl)-4-hydroxy-4-(3-trifluoromethyl-4-chlorophenyl)piperidine, 2j, (6.7 g) and triethylamine (4 mL) in dichloromethane (100 mL) was added dropwise at 0-5 °C a solution of chloroacetylchloride (2.6 mL) in dichloromethane (50 mL). The mixture was stirred overnight at room temperature. Water was added and pH was adjusted to >9. The organic phase was separated
30 and worked up as previously. The crude product was purified by column chromatography on silicagel (eluted with 4% triethylamine in ethyl acetate). Yield of pure 18a 6.2 g as an oil.

EXAMPLE 19

5-Chloro-1-[1-(6-methylaminocarbonylaminoindan-1-ylmethyl)piperidin-4-yl]-1H-indole, **19a**.

- 5 To a solution of 1-[1-(6-aminoindan-1-ylmethyl)piperidin-4-yl]-5-chloro-1H-indole, **10h** (0.9 g) in dichloromethane (10 mL) was added methylisocyanate (0.2 mL). The mixture was stirred at room temperature for 16 hours. Dichloromethane was evaporated. Upon addition of ethyl acetate the title compound **19a** crystallized. The crystalline product was filtered off and dried overnight at 80 °C *in vacuo*. Yield 0.6 g.
- 10 mp: 193-195 °C. ¹H NMR (DMSO-d₆) : δ 1.70-1.85 (m, 1H); 1.90-2.35 (m, 7H); 2.35-2.85 (m, 4H); 2.65 (d, 3H); 3.05-3.15 (m, 2H); 3.25 (quin, 1H); 4.35-4.45 (m, 1H); 5.95 (dt, 1H); 6.45 (d, 1H); 7.00-7.20 (m, 3H); 7.50-7.65 (m, 4H); 8.35 (s, 1H).

Pharmacological Testing

15

The compounds of the invention were tested in well recognized and reliable methods. The well-known 5-HT_{2A} antagonist MDL 100,907 and the well-known 5-HT_{1A} antagonist buspirone were included in the tests as reference compounds. The tests were as follows, and the results are given in the following Table 1.

20

Inhibition of ³H-8-OH-DPAT Binding to Serotonin 5-HT_{1A} Receptors in Rat Brain *in vitro*.

By this method the inhibition by drugs of the binding of the 5-HT_{1A} agonist ³H-8-OH-DPAT (1 nM) to 5-HT_{1A} receptors in membranes from rat brain minus cerebellum is determined *in vitro*. Accordingly, this is a test for affinity for the 5-HT_{1A} receptor. The test is performed as described by Hyttel et al., *Drug. Dev. Res.*, **1988**, 15, 389-404.

Inhibition of ³H-Ketanserin Binding to 5-HT₂ Receptors in Rat Cortex *in vitro*.

- 30 By this method the inhibition by drugs of the binding of ³H-Ketanserin (0,5 nM) to 5-HT_{2A} receptors in membranes from rat is determined *in vitro*. The method is described in Hyttel, *Pharmacology & Toxicology*, **61**, 126-129, 1987.

Table 1
Binding Data (IC₅₀ values in nM or % inhibition of binding at 100nM)

	Compound No	³ H 8-OH DPAT (5-HT _{1A})	³ H Ketanserin (5-HT _{2A})
	2a	480.	2.5
	2h	19%/100.	4.1
10	4a	11.	4.0
	4b	21.	2.9
	4c	28.	2.3
	4d	12.	5.0
	4e	38.	5.0
15	4f	84.	65.
	4g	31.	280.
	4h	11.	2.8
	4i	500.	2.1
	4j	45.	15.
20	4k	240	30.
	4l	26%/100	55.
	4m	120.	2.7
	4n	27.	3.9
	4o	11.	51.
25	4p	nt	nt
	5a	35	5.1
	5b	>100,000	37.
	5c	>1000.	3.9
	6a	1200	9.1
30	6b	1200	6.7
	6c	8200	9.9
	7a	13.	2.6
	11a	28.	42.
	11b	21.	44.
35	11c	16.	15.
	11d	29.	15.
	11e	27.	130.
	11f	8.0	21.
	11g	26.	180.
40	12a	340	3.5
	14a	120.	16.
	15a	91.	25%/100
	15b	25%/100	300.
	16a	170.	1.6
45	buspiron	41.	1300.
	MDL 100,907	nta	0.51

a) nt: not tested.

In addition to the above tests, the compounds of the invention were tested with respect to affinity for the dopamine D₂ receptor by determining their ability to inhibit the binding of ³H-spiroperidol to D₂ receptors by the method of Hyttel *et al*, *J. Neurochem.*, **1985**, *44*, 1615. Furthermore, they were tested with respect to their 5-HT reuptake inhibiting effect by measuring their ability to inhibit the uptake of ³H-serotonin in rat brain synapsomes *in vitro* by the method described by Hyttel and Larsen, *Acta Pharmacol. Tox.*, **1985**, *56*, suppl. 1, 146-153.

10 In general, the compounds of the invention have been found potently to inhibit both the binding of tritiated 8-hydroxy-2-dipropylaminotetralin (8-OH-DPAT) to 5-HT_{1A} receptors and the binding of ³H ketanserin to 5-HT_{2A} receptors *in vitro*. Some compounds only bind to one of the two serotonin receptor subtypes, 5-HT_{1A} or 5-HT_{2A}. In addition to these effects, a number of the compounds have proven to have 15 the further advantage of a potent 5-HT reuptake inhibiting effect. So, for example the compounds wherein R¹ is acetyl, R² is H and Ar is 3-indolyl substituted with halogen in the 6-position or 5-position or compounds wherein Ar is phenyl substituted with Cl in the 4-position show IC₅₀ values in the lower nanomolar range (1-65 nM).

20 Accordingly, the compounds of the invention are considered useful in the treatment of positive and negative symptoms of schizophrenia, other psychoses, anxiety disorders, such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder, depression, alcohol abuse, impulse control disorders aggression, side effects induced by conventional antipsychotic agents, ischaemic disease 25 states, migraine, senile dementia and cardiovascular disorders and in the improvement of sleep.

Formulation Examples

30

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art.

For example: Tablets may be prepared by mixing the active ingredient with ordinary

adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilization of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Typical examples of recipes for the formulation of the invention are as follows:

1) Tablets containing 5.0 mg of Compound **4a** calculated as the free base:

15	Compound 4a	5.0 mg
	Lactose	60 mg
	Maize starch	30 mg
	Hydroxypropylcellulose	2.4 mg
20	Microcrystalline cellulose	19.2 mg
	Croscarmellose Sodium Type A	2.4 mg
	Magnesium stearate	0.84 mg

2) Tablets containing 0.5 mg of Compound **4d** calculated as the free base:

25	Compound 4d	0.5 mg
	Lactose	46.9 mg
	Maize starch	23.5 mg
	Povidone	1.8 mg
	Microcrystalline cellulose	14.4 mg
30	Croscarmellose Sodium Type A	1.8 mg
	Magnesium stearate	0.63 mg


3) Syrup containing per millilitre:

41

	Compound 11f	25 mg
	Sorbitol	500 mg
	Hydroxypropylcellulose	15 mg
	Glycerol	50 mg
5	Methyl-paraben	1 mg
	Propyl-paraben	0.1 mg
	Ethanol	0.005 ml
	Flavour	0.05 mg
	Saccharin natrium	0.5 mg
10	Water	ad 1 ml
4)	Solution for injection containing per millilitre:	
	Compound 4a	0.5 mg
	Sorbitol	5.1 mg
15	Acetic acid	0.08 mg
	Water for injection	ad 1 ml

CLAIMS

1. A 4-aryl-1-(indanmethyl, dihydrobenzofuranmethyl or dihydrobenzothiophene-
5 methyl)piperidine, -tetrahydropyridine or -piperazine compound of general Formula
I



- 10 wherein one of X and Y is CH₂ and the other one is selected from the group
consisting of CH₂, O, and S;

the dotted line, emanating from Z, indicates an optional bond;

- when it does not indicate a bond Z is N, CH or COH; and when it indicates a bond Z
15 is C;

- Ar is phenyl, 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-pyrimidyl, 1-indolyl, 2-indolyl,
3-indolyl, 1-indol-2-onyl, 3-indol-2-onyl, 2- or 3-benzofuranyl, 2- or 3-benzothiophe-
nyl, 1-naphthyl or 2-naphthyl, each optionally substituted with halogen, lower alkyl,
20 lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl,
trifluoromethylsulfonyloxy, cycloalkyl, cycloalkyl-lower-alkyl, nitro, amino, lower
alkylamino, di-lower alkylamino, acylamino or C₁₋₂ alkylenedioxy;

- R₁ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl, cycloalk(en)yl, cycloalk(en)-
25 yl-lower alk(en/yn)yl, aryl, aryl-lower alkyl, acyl, thioacyl, lower alkylsulfonyl, trifluoro-
methylsulfonyl, arylsulfonyl,


R₁ is a group R⁹VCO- where V is O or S and R⁹ is lower alkyl, cycloalkyl, cycloalkyl-
lower-alkyl or aryl, or

R₁ is a group R¹⁰R¹¹NCO- or R¹⁰R¹¹NCS- wherein R¹⁰ and R¹¹ are independently

hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl, or R¹⁰ and R¹¹ together with the N-atom to which they are linked, form a pyrrolidinyl, piperidinyl or perhydroazepin group;

- 5 R² is hydrogen, lower alkyl, cycloalkyl or cycloalkyl-lower-alkyl;

or R¹ and R² together with the N-atom to which they are linked form a group ,



wherein Q is C=O, C=S or CH₂; T is NH, S, O or CH₂; and m is 1-4, inclusive;

10

R³ - R⁵ are independently hydrogen, halogen, lower alkyl, lower alkylcarbonyl, phenylcarbonyl, halogen substituted phenylcarbonyl, lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl, cycloalkyl, cycloalkyl-lower-alkyl or nitro;

15

R⁶ and R⁷ are each hydrogen or lower alkyl or they are linked together to constitute a 3 - 7-membered carbocyclic ring;

R⁸ is hydrogen or lower alkyl;

20

any alkyl, cycloalkyl or cycloalkylalkyl group present being optionally substituted with one or two hydroxy groups, which again are optionally esterified with an aliphatic or aromatic carboxylic acid; and any aryl substituent present being optionally substituted with halogen, lower alkyl, lower alkoxy, lower alkylthio, hydroxy, lower alkylsulfonyl, cyano, trifluoromethyl, trifluoromethylsulfonyloxy, cycloalkyl, cycloalkyl-lower-alkyl or nitro;

25

or a pharmaceutically acceptable acid addition salt thereof.

- 30 2. A compound of Claim 1, characterized in that X is CH₂ or S and Y is CH₂.

3. A compound of Claim 1 or 2, characterized in that R¹ is acyl, lower alkyl,

lower alkoxy, a group $R^{10}R^{11}NCO-$ or $R^{10}R^{11}NCS-$ wherein R^{10} is hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower-alkyl or aryl and R^{11} is hydrogen or lower alkyl or R^{10} and R^{11} together with the N-atom to which they are linked, form a pyrrolidinyl, piperidinyl or perhydroazepin group.

5

4. A compound of Claim 3, **characterized in that** R^1 is formyl, acetyl, methylaminocarbonyl, methylaminothiocarbonyl, dimethylaminocarbonyl, dimethylaminothiocarbonyl, methylsulfonyl, aminocarbonyl, cyclopropylcarbonyl, methyl, pyrrolidinylcarbonyl or 4-fluorophenylaminocarbonyl.

10

5. A compound of any of Claims 1 - 4, **characterized in that** R^2 is hydrogen or lower alkyl.

6. A compound of Claim 1, or 2, **characterized in that** R^1 and R^2 are linked together to form a 5-7 membered unsubstituted lactam ring or a pyrrolidinyl, piperidinyl or perhydroazepin.

20

7. A compound of any of Claims 1 - 6, **characterized in that** R^3-R^5 are hydrogen, fluoro, chloro, bromo, methyl, trifluoromethyl or acetyl and R^6-R^8 are all hydrogen.

8. A compound of any of Claims 1 - 4, **characterized in that** Ar is phenyl, 3-indolyl, 1-indolyl, or pyrimidyl or phenyl, 3-indolyl, 1-indolyl or pyrimidyl substituted with halogen.

25 9. A compound of any of Claims 1 - 5 and 7 - 8, **characterized in that** R^1 is acetyl, R^2 is H and Ar is indolyl or phenyl substituted with halogen.

10. A compound of Claim 9, **characterized in that** Ar is 3-indolyl substituted in the 5- or 6-position with chloro or fluoro or phenyl substituted in the 4-position with chloro.

30 11. A pharmaceutical composition **characterized in that** it comprises at least one compound of any of Claims 1-10 in a therapeutically effective amount and in

combination with one or more pharmaceutically acceptable carriers or diluents.

12. Use of a compound of any of Claims 1-10 or a pharmaceutically acceptable acid addition salt or prodrug thereof for the manufacture of a pharmaceutical preparation for the treatment of positive and negative symptoms of schizophrenia, other psychoses, anxiety disorders, generalised anxiety disorder, panic disorder, obsessive compulsive disorder, depression, alcohol abuse, impulse control disorders aggression, side effects induced by conventional antipsychotic agents, ischaemic disease states, migraine, senile dementia and cardiovascular disorders and for the improvement of sleep.

1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 95/00230

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 211/18, C07D 409/06, C07D 211/70, C07D 295/135, C07D 317/58,
C07D 211/52, C07D 211/10, C07D 401/12, 409/14, 401/04, 401/10, A61K 31/445,
According to International Patent Classification (IPC) or to both national classification and IPC 31/50

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0490772 A1 (ADIR ET COMPAGNIE), 17 June 1992 (17.06.92) --	1-12
A	EP 0428437 A1 (ADIR ET COMPAGNIE), 22 May 1991 (22.05.91) --	1-12
A	Derwent's abstract, No 93- 83841/10, week 9310, ABSTRACT OF SU, 845435-A1 (A Med Pharmacology Res Inst), 23 April 1992 (23.04.92) -- -----	1-12

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document but published on or after the international filing date	"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 11 Sept 1995	Date of mailing of the international search report 25-09-1995
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer Göran Karlsson Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT
Information on patent family members

28/08/95

International application No.
PCT/DK 95/00230

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A1- 0490772	17/06/92	AT-T- 125537	15/08/95
		AU-B- 638368	24/06/93
		AU-A- 8976291	18/06/92
		DE-D- 69111582	00/00/00
		FR-A,B- 2670491	19/06/92
		JP-A- 6025217	01/02/94
		JP-B- 7035377	19/04/95
		US-A- 5194437	16/03/93
EP-A1- 0428437	22/05/91	SE-T3- 0428437	
		AT-T- 110075	15/09/94
		AU-B- 631466	26/11/92
		AU-A- 6581990	16/05/91
		CA-A- 2029372	08/05/91
		DE-D,T- 69011628	30/03/95
		ES-T- 2062458	16/12/94
		FR-A,B- 2654104	10/05/91
		JP-A- 3188077	16/08/91
		JP-B- 6092403	16/11/94
		OA-A- 9468	15/11/92
		US-A- 5100902	31/03/92
		US-A- 5134147	28/07/92